Food safety is a key concern to us - this brochure is brought to you by:



TURKEY FARMERS OF CANADA LES ÉLEVEURS DE DINDON DU CANADA

turkeyfarmersofcanada.ca

1-888-245-8180 or ehfc.ca

EHFC



Canadian Partnership for Consumer Food Safety Education canfightbac.org



Design and production: Avalon Design & Marketing Inc. JUNE09 151500





Food Safety at **Home**



Your guide to safe food handling

More questions about safe food handling? Visit www.canfightbac.org or call your local public health office as listed in the blue pages of your phone book.

Home food safety the information you're hungry for...



Mishandling food in the home can result in foodborne illness. Foodborne illness results from eating foods that contain enough harmful microorganisms or their toxins to cause illness. Foodborne illness is often mistaken as a viral illness, like the 24-hour flu. Health Canada estimates that there are 11 to 13 million cases of foodborne illness in Canada each year. **The Facts...**

• For those at high risk - infants and young children, pregnant women, older adults and people with weakened immune systems - foodborne illness can result in very serious health problems, such as kidney failure.

• All foods, including fruits and vegetables, have the potential to cause illness since microorganisms can exist on all surfaces.

• Two out of three households in North America put themselves at risk for foodborne illness by not practising basic food safety steps at home.

You're in control...

There are **4 EASY steps** you can take at home to eliminate harmful bacteria and greatly reduce the risk of foodborne illness for your family.

- 1. Clean wash hands and surfaces often
- 2. Chill refrigerate or freeze foods promptly
- 3. **Separate** keep raw meat/poultry/seafood and their juices separate from one another and other foods
- 4. **Cook** cook foods to proper temperatures

Studies estimate that about eighty-five percent of all cases of foodborne illness could be prevented if food is handled properly. There are 4 simple safe food-handling steps you can take at home.



Clean: Wash hands and surfaces often. Proper hand washing may eliminate nearly half of all cases of foodborne illness.

Thoroughly wash hands using warm, soapy water.



• Wash your hands before you begin cooking and after you touch pets, change a diaper, cough or sneeze, use the phone, sort dirty laundry, use the restroom, take out the garbage or handle dirty dishes.

• Wash your hands for 20 seconds – that's two choruses of "Happy Birthday". Keep a nail scrub brush handy to get under your fingernails. Use a clean cloth or paper towels to dry.



• Mix 1 tsp (5 mL) bleach with 3 cups (750 mL) water to sanitize drains, cutting boards, taps, sinks, counter tops, sponges, pot scrubbers and fridge door handles. Store your bleach solution in a labelled spray bottle.

• Wash sponges with hot soapy water after every use. Change sponges, dishcloths, aprons and towels **often** machine wash in hot water. Replace sponges every few weeks.

• Wash all fruits and vegetables, including those that you peel or cut, like melons, oranges or cucumbers.

• Washing eggs is **not recommended** as this will remove their natural protective coating.

• Washing raw meat/poultry/seafood before cooking is **not recommended** as this can spread bacteria to kitchen surfaces and other foods.

Frequently sanitize your kitchen using a chlorine bleach solution.



Chill: Refrigerate/freeze foods promptly. Cold temperatures can prevent most harmful bacteria from growing.



Refrigerate perishables at less than 40°F (4°C). • Use the 2 Hour Rule in your home and while shopping – refrigerate or freeze all perishables within 2 hours of purchase or preparation. If the weather is hot, (greater than 80°F/26°C), reduce that time to 1 hour and use a cooler for perishables.

 Set your fridge temperature to be less than 40°F (4°C) and just slightly above the point where your lettuce will freeze. Check the temperature with a refrigerator thermometer. Avoid packing your fridge - cold air must be able to circulate.

• Store eggs in the main body of the fridge, not the door. This keeps eggs at a cooler, more consistent temperature.

• **Marinate foods in the fridge.** (If using a 30 minute quick-marinade product, follow package instructions).

Never defrost foods at room temperature.

 Meat/poultry/seafood defrosted in the microwave should be cooked immediately.

Cook thawed meat/poultry/seafood before refreezing.
If ice crystals are still present, it can be refrozen.

• Remove cooked poultry from bones of whole birds before storing in fridge or freezer.

• Divide leftovers into small portions to get foods cold faster. Refrigerate leftovers promptly and cover when cooled. Don't keep leftovers longer than 2 to 3 days.

Storage Chart

(from "packaged on" date or purchase date from butcher)

	Fridge	Freezer
Ground meats†/poultry, fresh seafood	1 day	2-3 months
Variety meats (e.g. liver, kidney, heart)	1-2 days	3-4 months
Stew meat, ribs, stir-fry strips, kabobs	2 days	3-6 months
Whole chicken/turkey or pieces	2-3 days	pieces 6 months whole 12 months
Steaks, roasts, chops	3 days	8-12 months
Cooked meats/poultry, cold cuts (open pkg)	3-4 days	2-3 months
Vacuum packed roasts/steaks (unopened)	See pkg date	10-12 months
Fresh shell eggs	See best before date	For egg freezing instructions visit eggs.ca

⁺For ground meats with longer storage time look for packs that have a 'best before date.' Once opened, use or freeze within one day.

You can't tell if food is safe by smelling or looking at it. IF IN DOUBT, THROW IT OUT.

Refrigerator defrosting is the safest way to thaw.

MIL

Separate: Don't cross-contaminate. Keep raw meat/poultry/seafood and their juices separate from one another and other foods during storage and preparation.



Keep foods covered.

• When shopping, keep packages of meat/poultry/seafood in bags separate from other groceries.

• Prevent meat/poultry/seafood juices from dripping onto other foods in the fridge - place on a plate and store on the bottom shelf.

• Place washed produce in clean containers, not back into their original unwashed packaging.

- Store eggs in their original carton to prevent them from absorbing odours and flavours from other foods.
- Wash scissors or blades used to cut open food packages.
- Ideally use two cutting boards: one for raw meat/ poultry/seafood; the other for cooked foods and washed fruits/vegetables.

• **Clean and sanitize cutting boards** using hot soapy water followed by a mild bleach solution (1 tsp/5 mL bleach: 3 cups/750 mL water); rinse. Plastic cutting boards can be cleaned and sanitized in the dishwasher.

• When checking cooked foods for seasoning, don't "double dip" - once you've done your tasting, don't put the tasting spoon back into the pot.

• When basting, brush sauce on cooked surfaces only. Be careful not to add sauce with a brush previously used on raw meat/poultry/seafood.

• Once marinade has been in contact with raw meat discard or bring to a rolling boil for 1 minute before using as a basting or dipping sauce. Do not reuse.

• Remove stuffing from poultry immediately after roasting or cook stuffing separately.

Use one plate or platter for raw meat/poultry/seafood and another for cooked.

Cook: Cook meat, poultry and eggs to proper temperatures.

Don't rely on the

'colour test' to

meats are done

(especially burgers

and meatloaves).

know when

• Use a food thermometer or temperature indicator. This is the ONLY way to tell if your food has reached a high enough internal temperature to destroy harmful bacteria.

• To know when burgers are done, check patty temperatures with a proper thermometer. Research shows that the colour of cooked ground beef can vary - beef burger patties may be brown in the centre before being cooked to a safe temperature.

• Enjoy steaks and roasts at medium-rare to medium doneness. Bacteria usually exist on the surface of these whole muscle cuts and are eliminated once the outside of the meat is well cooked. Note: Rolled steaks/roasts and fresh meats labelled "seasoned" should be cooked to medium doneness.

• Don't bring meat/poultry/seafood to room temperature before cooking.

• Finish cooking partially cooked meat/poultry/seafood immediately - do not hold to finish cooking at a later time.

- Serve cooked eggs or egg-rich foods (like rice pudding) immediately after cooking or refrigerate promptly and serve within 2 to 3 days.
- Stuff poultry just before cooking, stuffing loosely no more than two-thirds full.

• Use the Temperature Rules chart on next page, for safe doneness temperatures regardless of thermometer manufacturer's guidelines.

Temperature Rules for Safe Doneness

Ground beef/pork/veal/lamb	160°F (71°C)	
Ground chicken/turkey	165°F (74°C)	
Beef, lamb and veal roast and steaks	145°F (63°C) Med-rare 160°F (71°C) Medium 170°F (77°C) Well	
Pork chops/roasts/fresh cured ham	160°F (71°C) Medium	
Cured ham, fully cooked	Ready to eat	
Cured ham, partially cooked	160°F (71°C)	
Whole turkey (stuffed) and chicken (stuffed or not)	185°F (85°C)	
Stuffing (alone or in bird cavity)	165°F (74°C)	
Whole turkey (without stuffing)	185°F (85°C)	
Chicken/turkey pieces	165°F (74°C)	
Rolled stuffed beef roasts or steaks (e.g. London Broil)	160°F (71°C)	
Minute Steak (or meat labelled Delicatized/ Diced/Tenderized or Cubed Steak. NOT Fast-fry Steak)	160°F (71°C)	
Egg dishes, casseroles	165°F (74°C)	
Battered meat/seafood – Do not undercook. Cook following package directions		
Fresh meats marked "Seasoned" on label	160°F (71°C)	
Leftovers, reheated	165°F (74°C)	



The heat from proper cooking can kill bacteria.

Be a **GREAT** cook. Be a **SAFE** cook. Use a **food thermometer**.



Take the guesswork out of cooking -

Cook foods to **perfection**, so they are **tasty and juicy** – not overdone. **A food thermometer is an essential tool for a great cook.**

Keep your family safe -

Cook to **safe** internal temperatures. Be especially careful if you're cooking for those at high risk of serious illness from foodborne bacteria – infants/young children, pregnant women, older adults or those with chronic illness. Using a food thermometer or temperature indicator is the ONLY reliable way to test for safe internal temperatures.

An investment in safety -

Buy a digital instant-read thermometer or thermometer fork, that gives a temperature reading, <u>not just a doneness range</u>. Look for digital instant-read thermometers at quality hardware stores, cookware stores and restaurant equipment suppliers.

Thermometer Know-How

Rules of Thumb:

1. Take temperatures of thin foods like burgers within 1 minute of removal from heat, larger cuts like roasts, after 5 to 10 minutes.

2. Insert thermometer stem/indicator into the thickest part of the food, away from bone, fat or gristle.

3. Leave thermometer in food for at least 30 seconds before reading temperature.

4. When food has an irregular shape, like some beef roasts, check the temperature in several places.

5. Always wash the thermometer stem thoroughly in hot, soapy water after each use!

6. Review manufacturers' guidelines specific to the use of your thermometer or temperature indicator.



A food thermometer is an essential tool for a great cook.





Always wash the tongs, plate and thermometer stem used in checking partially cooked meats/poultry before using again.

* Doneness ranges should match temperatures indicated in Temperature Rules Chart.